

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/00, 9/02, C12P 1/00, C07D 231/06, 231/08, C11D 7/46, 7/60, 7/42, 3/18, 3/20, 3/26, 3/28, 3/30		A1	(11) International Publication Number: WO 97/06244 (43) International Publication Date: 20 February 1997 (20.02.97)
(21) International Application Number: PCT/EP96/03333 (22) International Filing Date: 29 July 1996 (29.07.96) (30) Priority Data: 9516244.2 8 August 1995 (08.08.95) GB 9525992.5 20 December 1995 (20.12.95) GB (71) Applicant (for all designated States except US): CIBA SPECIALTY CHEMICALS HOLDING INC. [CH/CH]; Klybeckstrasse 141, CH-4057 Basel (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): ZING, Rudolf [CH/CH]; Alemannenstrasse 2, CH-4106 Therwil (CH). FLETCHER, Ian, John [GB/CH]; Bürgenstal 27, CH-4312 Magden (CH). FREIERMUTH, Beat [CH/CH]; Oetlingerstrasse 170, CH-4057 Basle (CH). HUBER, Klaus [DE/DE]; Edith-Stein-Strasse 7, D-79110 Freiburg (DE). (74) Common Representative: CIBA SPECIALTY CHEMICALS HOLDING INC.; Patentabteilung, Klybeckstrasse 141, CH-4057 Basel (CH).		(81) Designated States: AL, AU, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: METHOD FOR ENHANCING THE ACTIVITY OF AN ENZYME, COMPOUNDS SHOWING ENZYME ACTIVITY ENHANCEMENT, AND DETERGENTS CONTAINING SUCH COMPOUNDS			
<div style="text-align: center;"> <div style="position: absolute; left: 640px; top: 610px;">(1)</div> </div>			
(57) Abstract A method of enhancing the activity of an enzyme, comprises adding to the enzyme, as a mediator for enhancing the activity of the enzyme, a compound having formula (1) in which A is a (hetero)aromatic residue; X is an -N=S or an -N=O group; Y is an -SH or an -OH group; Z is a water-solubilising group; x is 1 or 2; y is 1, 2 or 3; and z is 0, 1, 2, 3 or 4. The enzyme of enhanced activity is useful, e.g., in the degradative bleaching of pulp for paper production; in the degradative bleaching of stains on textiles during washing; in the modification, breakdown or bleaching of lignin, especially during the production of paper pulp; in the degradation of organic material such as a dyestuff in waste water; and in the degradative bleaching of migrating dyes in a washing process, so that inhibition of transfer of such dyes during washing is effected.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

METHOD FOR ENHANCING THE ACTIVITY OF AN ENZYME, COMPOUNDS SHOWING ENZYME ACTIVITY ENHANCEMENT, AND DETERGENTS CONTAINING SUCH COMPOUNDS

The present invention relates to a method of enhancing the activity of an enzyme with an enhancing agent for the enzyme and to the use of the enzyme of enhanced activity in a degradation process.

The use of a peroxidase enzyme in a process for the degradation, e.g. by bleaching, of a wide variety of substrates is already known.

For example, the use is known of a peroxidase enzyme in the bleaching of pulp for paper production, in the bleaching of textiles during washing, in the inhibition of dye transfer during washing and in the modification, breakdown or bleaching of lignin, especially during the production of paper pulp. In these known processes, the peroxidase enzyme is conventionally used together with hydrogen peroxide or a compound capable of producing hydrogen peroxide *in situ*.

Bourbonnais *et al.*, in FEBS letters (1990) 267, No.1, 99-102, have noted that, in the presence of certain compounds, termed mediators, such as the reactive dye Remazole Blue® or 2,2'-azinobis(3-ethylbenzthiazoline-6-sulfonate) (ABTS), laccase enzymes (polyphenol oxidases) can oxidize not only phenolic but also non-phenolic lignin subunits.

Although ABTS exhibits an excellent performance as a mediator for the bleaching of undesired dyestuffs, e.g. in the inhibition of dye transfer during washing, it is itself a dyestuff and suffers from the disadvantage that, after effecting its bleaching action, residual amounts of ABTS always remain in the treatment system.

In WO 94/12621, it has been proposed to use, as enhancers of the activity of peroxidases, organic chemical compounds consisting of at least two aromatic rings, at least one of which rings is substituted by one or more nitrogen, oxygen or sulfur atoms, the rings being optionally fused.

As merely one of many preferred sub-groups of such aromatic compounds, WO 94/12621 proposes naphthalene compounds which are optionally substituted by a wide variety of


groups in all possible positions on the naphthalene nucleus. The disclosed substituent groups, however, do not include the nitroso group.

Moreover, in WO 94/29510, there is described a process for modifying, breaking down or bleaching lignin using oxidation catalysts and suitable oxidizing agents. The catalysts are used in combination with an aliphatic, cycloaliphatic, heterocyclic or aromatic compound containing NO-, NOH- or H-N(R)-OH groups (in which R is not defined). Within this wide scope of compounds, N-hydroxy heterocyclic compounds such as hydroxybenzotriazole appear to be preferred. No aromatic compounds containing a nitroso group are specifically mentioned.

1-Hydroxybenzotriazole is indeed an excellent mediator. It is probably not sufficiently economic, however, for use in bleaching applications. Moreover, during enzyme reaction involving 1-hydroxybenzotriazole, there is obtained the by-product benzotriazole, which has been found to be mutagenic.

Surprisingly, it has now been found that using, as a mediator in the degradation of a substrate in the presence of an oxidoreductase enzyme, a (hetero)aromatic compound containing both a hydroxy and a nitroso group, and/or the analogous sulfur compounds, excellent levels of substrate degradation are obtained.

Accordingly, the present invention provides, as a first aspect, a method for enhancing the activity of an enzyme, comprising the use of the enzyme in combination with a mediator compound having the formula:



in which A is a (hetero)aromatic residue; X is an -N=S or, preferably, an -N=O group; Y is an -SH or, preferably, an -OH group; Z is a water-solubilising group, preferably a group -SO₂NH₂, -CO₂M or -SO₃M in which M is hydrogen or a cation; x is 1 or 2, preferably 1; y is 1, 2 or 3, preferably 1; and z is 0, 1, 2, 3 or 4, preferably 0, 1 or 2.

The mediator compound of formula (1) may be present in any of its possible tautomeric forms and/or transition metal complex forms. When present as a transition metal complex, the mediator compound of formula (1) is preferably in the form of an iron, manganese or cobalt metal complex. Moreover, in addition to the essential substituents X and Y, the residue A may also contain one or more further substituents such as C₁-C₄alkyl, C₁-C₄alkoxy, halogen and nitro groups.

The enzyme component, the activity of which is to be enhanced according to the present invention, may be any oxidoreductase or any fraction or synthetic mimic of such an enzyme. By enzyme mimics are meant compounds which simulate the prosthetic group of an oxidoreductase. Examples of oxidoreductases include oxidases, peroxidases, lignin peroxidases, manganese peroxidases and, in particular, laccases and laccase related enzymes.

The peroxidase enzyme may be any of those comprised by the enzyme classification EC 1.11.1.7, or any peroxidase activity-exhibiting fragment derived from such enzyme. Also of interest are synthetic or semi-synthetic derivatives of such enzymes, such as those comprising porphyrin ring systems or microperoxidases which are described, e.g., in US-4077768, EP-537381, WO 91/05858 or WO 92/16634. Other relevant enzymes are peroxidase-active fragments derived from cytochromes, hemoglobin or peroxidase enzymes, or synthetic or semi-synthetic derivatives thereof such as iron porphins, iron porphyrins, iron phthalocyanine and derivatives thereof. Such enzymes may be derived from animal, vegetable or microbial sources.

The laccase or related laccase enzyme may be any of those comprised by the enzyme classification EC 1.10.3.2, any catechol oxidase enzyme comprised by the enzyme classification EC 1.10.3.1, any bilirubin oxidase enzyme comprised by the enzyme classification EC 1.3.3.5 or any monophenol monooxygenase enzyme comprised by the enzyme classification EC 1.14.99.1.


The laccase or related laccase enzyme may also be one which can be produced by cultivation of a host cell transformed with a recombinant DNA vector carrying a DNA sequence encoding the laccase and carrying DNA sequences encoding functions allowing

the expression of the DNA sequence encoding the laccase. The cultivation may be conducted in a culture medium under conditions which permit the expression of the laccase enzyme. Finally, the laccase may be recovered from the culture medium.

The laccases are preferably produced from white rotting fungi, especially *Phanerochaete chrysosporium* and *Coriolus (Trametes) versicolor*.


In the mediator compounds of formula (1), A may be a purely aromatic residue, preferably a phenyl residue or, especially, a naphthyl residue, or it may be a heteroaromatic residue, that is an aromatic residue containing one or more heteroatoms, preferably nitrogen, oxygen or sulfur atoms, such a pyrimidine or coumarin residue, or an aromatic residue substituted with a heterocyclic residue. Both aromatic and heteroaromatic residues A include mono- and polycyclic residues. In the case of the latter, the individual rings may be fused together, or linked together via a bridging member such as a direct bond, $-(CH_2)_n-$, $-(CH=CH)_n-$, $-N=N-$, $-O-$, $-CO-$, $-NH-$, $-N=N-$, $-S-$ or $-SO_2-$ in which n is an integer from 1 to 6.

Examples of preferred purely aromatic mediator compounds of formula (1) are those having the formula:




in which Z, x, y and z have their previous significance.

Preferred purely aromatic compounds of formula (2) are those having the formula:




or the formula:




in which Z and z have their previous significance.

Further examples of preferred purely aromatic mediator compounds of formula (1) are those having the formula:




in which Z, x, y and z have their previous significance, Me is a transition metal and p is the valency of the transition metal Me.

Examples of preferred heteroaromatic mediator compounds of formula (1) are pyrimidine derivatives having the formula:




(the commercially available compound violuric acid), which can also exist in other tautomeric forms such as that having the formula:


- 6 -




or having the formula:



(1,3-dimethyl-5-nitrosobarbituric acid; Beilstein Ell, vol. 24, 307); and coumarin derivatives such as that having the formula:




(4-hydroxy-3-nitrosocoumarin; Beilstein E III/IV, vol. 17, 6737); as well as compounds having the formula:



(US-A-2 646 409, Example 19); or having the formula:

- 7 -





(having m.pt 230°C.(decomp.) and the elemental analysis (calculated for $C_{10}H_{10}N_4SO_3$) C=42.85; H=3.57; N=19.85; S=11.36; O=22.67; found C=42.7; H=3.5; N=19.8; S=11.3; O=22.7 and produced analogously to the procedure described in US-A-2 646 409, Example 19).




In the compounds of formula (1), (2), (3) or (4), when M is a cation, it is preferably an alkali metal cation, especially a sodium cation, or a cation formed from an amine.

When, in the compounds of formula (1), (2), (3) or (4), Z is a water-solubilising group, then it will be understood that the compounds of formula (1), (2), (3) or (4) may be used in the form of any desired hydrate.


Particularly preferred purely aromatic mediator compounds of formula (1) include those having one of the formulae:



- 8 -





(Beilstein E II, vol. 11, page 150).



(C.I.10020; Acid Green B; Naphthol Green B); which may be in the tautomeric form having the formula:

- 9 -



(Berichte Deutschen Chem. Ges. 42, page 4266).

The mediator compounds are mostly known compounds and can be obtained commercially or by methods which are well known. The compound of formula (10), however, is a new compound and, as such, forms a second aspect of the present invention.

The present invention provides, as a third aspect, a process for the degradation of a substrate comprising contacting the substrate with a bleach liquor comprising an oxidizing agent, an enzyme and a mediator for the enzyme, according to the first aspect of the present invention.

If a peroxidase enzyme is used, the oxidizing agent used in the degradation process according to the present invention is preferably hydrogen peroxide or a compound capable of producing hydrogen peroxide *in situ*, such as a perborate, a percarbonate, a peroxycarboxylic acid or a salt thereof.

On the other hand, if a laccase or laccase-related enzyme is used, the oxidizing agent used in the degradation process according to the present invention is preferably oxygen, conveniently supplied as air or pure oxygen, optionally with the application of pressure.

In addition to the above-mentioned enzymes, the degradation process of the invention may also include the addition of hemicellulases, cellulases, pectinases, amylases and lipases to the reaction mixture. Mixtures of two or more enzymes may also be used and, under certain circumstances, synergistic effects may be observed with the use of such mixtures of enzymes.

The oxidizing agent is preferably added at the start of the degradation process and preferably in an amount of from 0.0001 to 5.0% by weight, more preferably from 0.01 to 1.0% by weight, based on the total weight of the substrate to be degraded.

The degradation process according to the present invention is preferably applied to the degradative bleaching of pulp for paper production; in the degradative bleaching of stains on a) textiles during washing thereof with laundry detergents or on b) dishes during washing thereof with dishwashing detergents; in the modification, breakdown or bleaching of lignin, especially during the production of paper pulp; in the degradation of organic material such as a dyestuff in waste water; and in the degradative bleaching of migrating dyes in a washing process, so that inhibition of transfer of such dyes during washing is effected.

When the process of the present invention is applied to degradative bleaching of pulp, the process is generally conveniently operated at a temperature in the range of from 25-80°C., preferably at a temperature in the range of from 40-60°C. The process is preferably conducted in the presence of air or oxygen and at a pressure ranging from normal pressure up to 10 bar excess pressure. The consistency of the process reaction mixture preferably ranges from 0.5-40%.

The process of the present invention may be applied to the production of pulps in general. Thus, the process may be used for the delignification (bleaching) of sulfate-, sulfite-, organosolve- and wood pulp.

The present invention further provides a method for the degradative bleaching of stains on a) textiles during washing thereof with laundry detergents or on b) dishes during washing thereof with dishwashing detergents, comprising washing the textiles or dishes with a

laundry or dishwashing detergent which contains an enzyme having enhanced activity by the addition of a mediator compound of formula (1) according to the present invention.

The present invention also provides a method for inhibiting the re-absorption of migrating dyes in the wash liquor, comprising introducing into a wash liquor containing a peroxide detergent, from 0.001 to 150 mg, per litre of wash liquor, of an enzyme having enhanced activity by the addition of a mediator compound of formula (1) according to the present invention.

When the method of the present invention is applied to the degradative bleaching of stains on textiles during washing thereof with laundry detergents, or for inhibiting the re-absorption of migrating dyes in the wash liquor, comprising introducing into a wash liquor containing a peroxide detergent, the method is generally conveniently operated at a temperature in the range of from 15-80°C., preferably at a temperature in the range of from 25-50°C.

The present invention further provides a detergent composition comprising:

- i) 5-90%, preferably 5-70% of A) an anionic surfactant and/or B) a nonionic surfactant;
- ii) 5-70%, preferably 5-50%, especially 5-40% of C) a builder;
- iii) 0.01-30%, preferably 1-12% of D) a peroxide; and
- iv) 0.001-2%, preferably 0.02-2% of E) an enzyme having enhanced activity by the addition of a mediator compound of formula (1) according to the present invention, each by weight, based on the total weight of the detergent.

Each of components A) to E) may be a single compound or a mixture of compounds.

The detergent may be formulated as a solid; or as a non-aqueous liquid detergent, containing not more than 5, preferably 0-1 wt.% of water, and based on a suspension of a builder in a non-ionic surfactant, as described, e.g., in GB-A-2158454.

Preferably, the detergent is in powder or granulate form.

Such powder or granulate forms may be produced by firstly forming a base powder by spray-drying an aqueous slurry containing all the said components, apart from the components D) and E); then adding the components D) and E) by dry-blending them into

the base powder. In a further process, the component E) may be added to an aqueous slurry containing components A), B) and C), followed by spray-drying the slurry prior to dry-blending component D) into the mixture. In a still further process, component B) is not present, or is only partly present in an aqueous slurry containing components A) and C); component E) is incorporated into component B), which is then added to the spray-dried base powder; and finally component D) is dry-blended into the mixture.

The anionic surfactant component A) may be, e.g., a sulphate, sulphonate or carboxylate surfactant, or a mixture of these.

Preferred sulphates are alkyl sulphates having 12-22 carbon atoms in the alkyl radical, optionally in combination with alkyl ethoxy sulphates having 10-20 carbon atoms in the alkyl radical.

Preferred sulphonates include alkyl benzene sulphonates having 9-15 carbon atoms in the alkyl radical.

In each case, the cation is preferably an alkali metal, especially sodium.

Preferred carboxylates are alkali metal sarcosinates of formula $R-CO(R^1)CH_2COOM^1$ in which R is alkyl or alkenyl having 9-17 carbon atoms in the alkyl or alkenyl radical, R^1 is C_1-C_4 alkyl and M^1 is alkali metal.

The nonionic surfactant component B) may be, e.g., a condensate of ethylene oxide with a C_9-C_{15} primary alcohol having 3-8 moles of ethylene oxide per mole.

The builder component C) may be an alkali metal phosphate, especially a tripolyphosphate; a carbonate or bicarbonate, especially the sodium salts thereof; a silicate; an aluminosilicate; a polycarboxylate; a polycarboxylic acid; an organic phosphonate; or an aminoalkylene poly (alkylene phosphonate); or a mixture of these.

Preferred silicates are crystalline layered sodium silicates of the formula $\text{NaHSi}_m\text{O}_{2m+1}$ or $\text{Na}_2\text{Si}_m\text{O}_{2m+1} \cdot p\text{H}_2\text{O}$ in which m is a number from 1.9 to 4 and p is 0 to 20.

Preferred aluminosilicates are the commercially-available synthetic materials designated as Zeolites A, B, X, and HS, or mixtures of these. Zeolite A is preferred.

Preferred polycarboxylates include hydroxypolycarboxylates, in particular citrates, polyacrylates and their copolymers with maleic anhydride.

Preferred polycarboxylic acids include nitrilotriacetic acid and ethylene diamine tetra-acetic acid.

Preferred organic phosphonates or aminoalkylene poly (alkylene phosphonates) are alkali metal ethane 1-hydroxy diphosphonates, nitrilo trimethylene phosphonates, ethylene diamine tetra methylene phosphonates and diethylene triamine penta methylene phosphonates.

The peroxide component D) may be any organic or inorganic peroxide compound, described in the literature or available on the market, which bleaches textiles at conventional washing temperatures, e.g. temperatures in the range of from 10°C. to 90°C. In particular, the organic peroxides are, for example, monoperoxides or polyperoxides having alkyl chains of at least 3, preferably 6 to 20, carbon atoms; in particular diperoxydicarboxylates having 6 to 12 C atoms, such as diperoxyperazates, diperoxypersebacates, diperoxyphthalates and/or diperoxydodecanedioates, especially their corresponding free acids, are of interest. It is preferred, however, to employ very active inorganic peroxides, such as persulphate, perborate and/or percarbonate. It is, of course, also possible to employ mixtures of organic and/or inorganic peroxides. Peroxides can have different crystalline forms and/or different degrees of hydration. They may be used in admixture with other organic or inorganic salts, thereby improving their stability to storage.

The addition of the peroxides to the detergent is effected, in particular, by mixing the components, for example by means of screw-metering systems and/or fluidized bed mixers.

The detergents may contain, in addition to the combination according to the invention, one or more of fluorescent whitening agents, such as a bis-triazinylamino-stilbene-disulphonic acid, a bis-triazolyl-stilbene-disulphonic acid, a bis-styryl-biphenyl, a bis-benzofuranyl-biphenyl, a bis-benzoxalyl derivative, a bis-benzimidazolyl derivative, a coumarine derivative or a pyrazoline derivative; soil suspending agents, for example sodium carboxymethylcellulose; salts for adjusting the pH, for example alkali or alkaline earth metal silicates; foam regulators, for example soap; salts for adjusting the spray drying and granulating properties, for example sodium sulphate; perfumes; and also, if appropriate, antistatic and softening agents; such as smectite clays; enzymes, such as amylases; photobleaching agents; pigments; and/or shading agents. These constituents should, of course, be stable to the bleaching system employed.

A particularly preferred detergent co-additive is a polymer known to be useful in preventing the transfer of labile dyes between fabrics during the washing cycle. Preferred examples of such polymers are polyvinyl pyrrolidones, optionally modified by the inclusion of an anionic or cationic substituent, especially those having a molecular weight in the range from 5000 to 60,000, in particular from 10,00 to 50,000. Preferably, such polymer is used in an amount ranging from 0.05 to 5%, preferably 0.2-1.7% by weight, based on the weight of the detergent.

The following Examples further illustrate the present invention.

Examples 1 to 3

The oxidation of C.I. Direct Blue 1 is monitored spectroscopically at 610 nm (the absorption peak of the dye) in a spectrophotometer.

The reagents are added, using a pipette, in the following sequence, to a cuvette having a layer thickness of 1 cm:

acetate buffer pH 5




C.I.Direct Blue 1

H₂O


mediator under test and

laccase enzyme (polyphenol oxidase produced from *Trametes versicolor*).

The test mediators used according to the present invention are those having the following formulae:



For the purpose of comparison, an experiment is conducted using, as mediator, the known mediator compound N-hydroxybenzotriazole having the formula:



Directly after the addition of the enzyme, the mixture is stirred with the tip of the pipette for about 15 seconds. 20 seconds after the addition of the enzyme, the spectroscopic measurement at 610 nm is commenced and the spectrum is monitored for 8 minutes. The temperature of the reagents and the cuvette is set at 25°C. Each experiment is conducted at pH 5. The volume of liquid in the cuvette is adjusted to 3 ml for each experiment by adding the necessary amount of water.

In the following Table 1, there are shown, for each reagent, the concentration of the reagent in the basic solution, the amount of the sample of reagent removed and the concentration of the reagent in the final solution.

Table 1

Reagent	Concentration of basic solution	Amount of sample taken	Final concentration in cuvette
acetate buffer pH 5	2.86 mole/l	25 µl	$20 \cdot 10^{-3}$ mole/l
Direct Blue 1	$7.54 \cdot 10^{-4}$ mole/l	100 µl	$2.51 \cdot 10^{-5}$ mole/l
mediator: Compound 101 Compound 103 Compound 104 HOBT	$1.42 \cdot 10^{-3}$ mole/l	21 µl	$10 \cdot 10^{-6}$ mole/l
laccase	6.0 U/ml	50 µl	0.1 U/ml

The concentration unit U (µmol/min) given for the laccase is defined in Bourbonnais *et al.*, in FEBS letters (1990) 267, No.1, 99-102 (supra).

The results of the dye oxidation tests, showing the optical density (OD) measurements at 0 and at 8 minutes, and the rate of increase (initial gradient) of the respective OD curves, are set out in the following Table 2.

Table 2

Mediator	OD _{0 min}	OD _{8 min}	Initial Gradient x 1000
none (laccase alone)	0.90	0.67	31.73
HOBT	0.90	0.56	54.18
Compound 101	0.83	0.40	231.17
Compound 103	0.90	0.36	79.72
Compound 104	0.89	0.33	108.98


The rate of dye oxidation obtained with the mediators used according to the invention is clearly greatly superior to that in the control experiment or using the known mediator HOBT.

Similar results are obtained when the compounds of formula (101), (103) and (104) are replaced by those of formula (105), (106), (107), (6), (7), (8), (9) or (10).

Example 4

1.0 ml of 1M H₂SO₄ is added to of moist softwood pulp (dry softwood pulp content 10.0 g) in order to maintain the pulp at an acidic pH.

0.05 g of a mediator compound of formula:



is dissolved in 20 mls of water. This mediator solution, together with 4 mls of a 0.5% by weight aqueous solution of EDTA and - mls of water are then added to the pulp in order to provide, in the mixture, a concentration of 5 mg of the mediator compound of formula (10) per g of pulp and a concentration of 2 mg of EDTA per g of pulp. Over a period of 60 seconds, the resulting mixture is thoroughly homogenized. To the homogenized mixture there are added Versicolor laccase (EC number 1.10.3.2), 1.9 mls of an aqueous 2.6M sodium acetate buffer solution (pH 5), 0.25 ml of a 20% by weight aqueous solution of MgSO_4 , to give a concentration of 3.5 U of laccase per g of pulp, a concentration of 0.5 mg of MgSO_4 per g of pulp and a concentration of 50 mM of sodium acetate in the resulting mixture. The mixture so obtained is kneaded for 60 seconds and the pH thereof is adjusted to 5, if necessary, by the addition of further 1M H_2SO_4 . The total weight of the mixture is then adjusted to 100 g by the addition of further water.

The kneaded mixture is placed in a steel container from which the air is then evacuated using a water jet pump. The container is then charged with oxygen to a pressure of 3 bar. The container is then heated at 50°C. for 225 minutes and subsequently the oxygen is removed using a water jet pump.

The reaction mixture is washed with 1 litre of warm (50°C.) water in a Buchner funnel; then extracted with a warm (70°C.) aqueous NaOH solution, using 2 g of NaOH per 100 g of pulp; and finally re-washed with 1 litre of warm (50°C.) water in a Buchner funnel. The total weight of the alkaline mixture (including 10 g of pulp) is then adjusted to 100 g by the addition of further water.

The Kappa number of the washed material is then determined according to the test method SCAN-C 1:77 of the Scandinavian Pulp and Paper Board. Compared with the initial Kappa number, a Kappa number reduction of 26% is obtained.

Example 5

Using a beaker laboratory dyeing apparatus having 50 mls containers, respective wash liquids are made up containing, in 25 mls of water, a) a borax buffer (pH 9) or a commercial detergent; b) a peroxidase from horse radish (Fluka No. 77333) dissolved in a phosphate buffer (pH 6); and c) the compound of formula (101). The wash liquor is pre-heated to 40°C. Test cotton swatches are immersed in the respective wash liquids and then 5 mls of an aqueous solution, also pre-heated to 40°C., of Cibacron Marine C-B dye and hydrogen peroxide are added.

The reagent concentrations are shown in the following Table 3.

Table 3

Reagent	Final concentration in wash liquor	
	buffer system	detergent system
borax buffer	20 mM	-
commercial detergent	-	4.4 g/l
H ₂ O ₂	200 µM	200 µM
compound (101) (stock solution 20:80 ethanol/water)	90 µM	90 µM
peroxidase, 727 U/mg	0.56 U/ml	0.56 U/ml
Cibacron Marine C-B	20 µM	8 µM
cotton swatches	2.8 g	3.75 g

The total volume in the wash container is 30 mls.

1U of Fluka No. 77333 corresponds to the amount of enzyme which oxidizes 1 μ mol of ABTS per minute at pH 6 and at 25°C.

During the washing process, the temperature is held at 40°C. for 30 minutes. The swatches are then removed from the containers, rinsed with water, dried and ironed. The brightness values (Y) of the washed samples, as well as those of control samples, are then determined spectrophotometrically. the results are set out in the following Table 4.


Table 4

Example	wash conditions	Y	
		buffer system	detergent system
-	control: cotton untreated	92.5	92.5
-	control: no enzyme or compound (101)	72.8	82.0
-	control: enzyme but no compound (101)	74.8	82.3
5	complete system	85.4	90.0

The results in Table 4 demonstrate that the method of the present invention is very effective in inhibiting the transfer of the dye on to the cotton and thus maintains, to a great extent, the original brightness of the washed cotton. The results obtained compare favourably with those resulting from the use of a commercial detergent.

Claims

1. A method for enhancing the activity of an enzyme, comprising adding to the enzyme, as a mediator for enhancing the activity of the enzyme, a compound having the formula:



in which A is a (hetero)aromatic residue; X is an -N=S or an -N=O group; Y is an -SH or an -OH group; Z is a water-solubilising group; x is 1 or 2; y is 1, 2 or 3; and z is 0, 1, 2, 3 or 4; or a tautomer or transition metal complex of a compound of formula (1).

2. A method according to claim 1 in which X is an -N=O group.

3. A method according to claim 1 or 2 in which Y is an -OH group.

4. A method according to any of the preceding claims in which Z is a group -SO₂NH₂, -CO₂M or -SO₃M in which M is hydrogen or a cation.

5. A method according to any of the preceding claims in which x is 1.


6. A method according to any of the preceding claims in which y is 1.

7. A method according to any of the preceding claims in which z is 0, 1 or 2.

8. A method according to any of the preceding claims in which the enzyme component is an oxidoreductase or any fraction or synthetic mimic of such an enzyme.


9. A method according to any of the preceding claims in which the oxidoreductase is an oxidase, peroxidase, lignin peroxidase, manganese peroxidase or laccase.

10. A method according to claim 9 in which the oxidoreductase is a laccase or laccase related enzyme.
11. A method according to any of the preceding claims in which a compound of formula (1) is used in which A is an aromatic residue.
12. A method according to claim 11 in which A is a naphthyl residue.
13. A method according to claim 12 in which the compound of formula (1) has the formula:




in which Z, x, y and z are as defined in claim 1.

14. A method according to claim 13 in which the compound of formula (2) has the formula:








or the formula:




in which Z and z are as defined in claim 1.


15. A method according to claim 14 in which the compound of formula (1) has one of the formulae:



- 24 -




16. A method according to claim 12 in which the compound of formula (1) has the formula:




in which Z, x, y and z are as defined in claim 1, Me is a transition metal and p is the valency of the transition metal Me.

17. A method according to claim 16 in which the compound of formula (5) has the formula:




or the tautomeric form having the formula:






18. A method according to any of claims 1 to 10 in which a compound of formula (1) is used in which A is an heteroaromatic residue.



19. A method according to claim 18 in which the compound of formula (1) has one of the formulae:



or the tautomeric form having the formula:



- 26 -



20. A method according to any of the preceding claims in which, in the compounds of formula (1), (2), (3) or (4), when M is a cation, it is an alkali metal cation or a cation formed from an amine.

21. A method according to any of the preceding claims in which when, in the compounds of formula (1), (2), (3) or (4), Z is a water-solubilising group, then the compounds of formula (1), (2), (3) or (4) are used in the form of any desired hydrate.

22. A process for the degradation of a substrate comprising contacting the substrate with a bleach liquor comprising an oxidizing agent, an enzyme and a mediator compound of formula (101) for the enzyme as defined in any of claims 1 to 21.

23. A process according to claim 22 in which the oxidizing agent is added at the start of the process in an amount of from 0.0001 to 5.0% by weight, based on the total weight of the substrate.

24. A process according to claim 23 in which the oxidizing agent is added in an amount of from 0.01 to 1.0% by weight, based on the total weight of the substrate.

25. A process according to any of claims 22 to 24 in which a peroxidase enzyme is used and the oxidizing agent is hydrogen peroxide or a compound capable of producing hydrogen peroxide *in situ*.

26. A process according to claim 25 in which the oxidizing agent is a perborate, a percarbonate, a peroxycarboxylic acid or a salt thereof.

27. A process according to any of claims 22 to 24 in which a laccase enzyme is used and the oxidizing agent is oxygen and is optionally applied under pressure.

28. A process according to any of claims 22 to 27 in which the degradation process is applied to the degraditive bleaching of pulp for paper production.

29. A process according to any of claims 22 to 27 in which the degradation process is applied to the degraditive bleaching of stains on textiles during washing with laundry detergents or to the degraditive bleaching of stains on dishes during washing of the dishes with dishwashing detergents.

30. A process according to any of claims 22 to 27 in which the degradation process is applied to the modification, breakdown or bleaching of lignin.

31. A process according to claim 30 in which the degradation process is applied to the production of paper pulp.

32. A process according to any of claims 22 to 27 in which the degradation process is applied to the degradation of organic material such as a dyestuff in waste water.

33. A process according to claim 32 in which the organic material is a dyestuff.

34. A process for inhibiting the re-absorption of migrating dyes in the wash liquor, comprising introducing into a wash liquor containing a peroxide detergent, from 0.001 to 150 mg, per litre of wash liquor, of an enzyme having enhanced activity by virtue of the addition of a mediator compound according to any of claims 1 to 21.

35. A detergent composition comprising:

- i) 5-90% of A) an anionic surfactant and/or B) a nonionic surfactant;
- ii) 5-70% of C) a builder;
- iii) 0.01-30% of D) a peroxide; and
- iv) 0.001-2% of E) an enzyme having enhanced activity by the addition of a mediator compound according to any of claims 1 to 21, each by weight, based on the total weight of the detergent.

36. A detergent composition according to claim 35 comprising:

- i) 5-70% of A) an anionic surfactant and/or B) a nonionic surfactant;
- ii) 5-50% of C) a builder;
- iii) 1-12% of D) a peroxide; and
- iv) 0.001-2% of E) an enzyme having enhanced activity by the addition of a mediator compound according to any of claims 1 to 21, each by weight, based on the total weight of the detergent.

37. A composition according to claim 36 comprising:

- i) 5-70% of A) an anionic surfactant and/or B) a nonionic surfactant;
- ii) 5-40% of C) a builder;
- iii) 1-12% of D) a peroxide; and
- iv) 0.02-2% of E) an enzyme having enhanced activity by the addition of a mediator compound according to any of claims 1 to 21, each by weight, based on the total weight of the detergent.

38. A detergent composition according to any of claims 35 to 37 comprising 0.5-5% by weight of a polymer useful in preventing the transfer of labile dyes between fabrics during a washing cycle.

39. A detergent composition according to claim 38 comprising 0.2-1.7% of the polymer.


40. A detergent composition according to claim 38 or 39 in which the polymer is a polyvinylpyrrolidone optionally containing an anionic or cationic substituent.

41. A detergent composition according to any of claims 35 to 40 in which the detergent is in powder or granulate form.

42. A detergent composition according to any of claims 35 to 40 in which the detergent is in liquid form and contains 0-5% water.

43. A detergent composition according to claim 42 in which the detergent is in liquid form and contains 0-1% water.

44. A compound having the formula:



INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/EP 96/03333

A. CLASSIFICATION OF SUBJECT MATTER					
IPC 6	C12N9/00	C12N9/02	C12P1/00	C07D231/06	C07D231/08
	C11D7/46	C11D7/60	C11D7/42	C11D3/18	C11D3/20
	C11D3/26	C11D3/28	C11D3/30		
According to International Patent Classification (IPC) or to both national classification and IPC					
B. FIELDS SEARCHED					
Minimum documentation searched (classification system followed by classification symbols)					
IPC 6	C12N	C12P	C07D	C11D	
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched					
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)					
C. DOCUMENTS CONSIDERED TO BE RELEVANT					
Category *	Citation of document, with indication, where appropriate, of the relevant passages				Relevant to claim No.
A	ACTA CHEM. SCAND., SER. B, vol. B42, no. 6, 1988, pages 414-416, XP002018936 LAGERCRANTZ C.: "Oxidation of Some "-Oximino-1,3-dioxo Compounds to Iminoxy Radicals by Horseradish Peroxidase and Hydrogen Peroxide" see the whole document ---				1-44
A	WO,A,95 01426 (NOVO NORDISK A/S) 12 January 1995 see the whole document ---				1-44
A	EP,A,0 361 470 (FIJIREBIO INC.) 4 April 1990 see the whole document ---				1-44
-/--					
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.					
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family					
Date of the actual completion of the international search			Date of mailing of the international search report		
20 November 1996			06. 12. 96		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016			Authorized officer Douschan, K		

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/EP 96/03333

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,94 12620 (NOVO NORDISK A/S) 9 June 1994 see the whole document ---	1-44
A	WO,A,94 29510 (CALL H.-P.) 22 December 1994 cited in the application see the whole document ---	1-44
A	WO,A,94 12621 (NOVO NORDISK A/S) 9 June 1994 cited in the application see the whole document -----	1-44

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/EP 96/03333

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9501426	12-01-95	AU-A- 6924594	24-01-95
		BR-A- 9406868	26-03-96
		CA-A- 2165283	12-01-95
		CN-A- 1126490	10-07-96
		EP-A- 0707637	24-04-96
		FI-A- 956329	23-02-96

EP-A-0361470	04-04-90	DE-D- 68911633	03-02-94
		DE-T- 68911633	07-04-94
		JP-A- 2174694	06-07-90
		US-A- 5171668	15-12-92

WO-A-9412620	09-06-94	CA-A- 2150562	09-06-94
		CA-A- 2150563	09-06-94
		WO-A- 9412621	09-06-94
		EP-A- 0677102	18-10-95
		EP-A- 0679183	02-11-95
		FI-A- 952647	31-05-95
		FI-A- 952648	28-07-95
		JP-T- 8503371	16-04-96
		JP-T- 8506009	02-07-96

WO-A-9429510	22-12-94	AU-A- 7124094	03-01-95
		AU-A- 7739794	03-01-95
		BR-A- 9406854	26-03-96
		CA-A- 2165426	17-12-94
		CA-A- 2182182	22-12-94
		CN-A- 1127523	24-07-96
		CZ-A- 9503325	15-05-96
		WO-A- 9429425	22-12-94
		EP-A- 0739433	30-10-96
		EP-A- 0705327	10-04-96
		FI-A- 956023	25-01-96
		FI-A- 961157	13-03-96
		NO-A- 955111	07-02-96
		NO-A- 961205	25-03-96

WO-A-9412621	09-06-94	CA-A- 2150562	09-06-94
		EP-A- 0679183	02-11-95
		FI-A- 952648	28-07-95

INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/EP 96/03333

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9412621		JP-T- 8506009	02-07-96
		AU-A- 7937194	22-05-95
		CA-A- 2175047	04-05-95
		WO-A- 9511964	04-05-95
		EP-A- 0730641	11-09-96
		CA-A- 2150563	09-06-94
		WO-A- 9412620	09-06-94
		EP-A- 0677102	18-10-95
		FI-A- 952647	31-05-95
		JP-T- 8503371	16-04-96
